

## Sex differences in serotonin enhancement of capsaicin-evoked calcitonin gene-related peptide release from human dental pulp

Dayna R. Loyd<sup>a,\*</sup>, Xiaoling X. Sun<sup>a</sup>, Erin E. Locke<sup>a</sup>, Margaux M. Salas<sup>b</sup>, Kenneth M. Hargreaves<sup>a,b,c</sup>

<sup>a</sup> Department of Endodontics, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>b</sup> Department of Cellular and Structural Biology, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>c</sup> Department of Pharmacology, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

### ARTICLE INFO

#### Article history:

Received 8 May 2012

Received in revised form 11 June 2012

Accepted 18 June 2012

#### Keywords:

CGRP

5HT

Craniofacial

Orofacial

Teeth

Pain

### ABSTRACT

Serotonin (5HT) is a pronociceptive mediator in the periphery, and evidence implicates involvement in trigeminal pain processing. However, the mechanism(s) by which 5HT modulates trigeminal nociceptors remains unclear. Trigeminal pain can be evoked by the transient receptor potential V1 channel (TRPV1), which is expressed by nociceptive trigeminal neurons and induces release of proinflammatory calcitonin gene-related peptide (CGRP). In our preclinical models, 5HT evoked thermal hyperalgesia and enhanced calcium influx and CGRP release from the TRPV1 population of trigeminal nociceptors. Whether this occurs in humans is unknown. As dental pulp is densely innervated by trigeminal nociceptors, routine tooth extractions offer a unique opportunity to examine whether 5HT enhances CGRP release from human nociceptors. Pulpal tissue was collected from 140 extracted molar teeth from men and women, and basal release samples were collected before treatment with saline or 5HT 100  $\mu\text{mol/L}$ . CGRP release was then stimulated with the TRPV1 agonist capsaicin 1  $\mu\text{mol/L}$  and quantitated by enzyme immunoassay. Additional samples were collected for Western blots to examine 5HT receptor expression. We report that 5HT induced a significant increase in capsaicin-evoked CGRP release, and that this enhancement was observed only in female dental pulp, with no effect of 5HT on male dental pulp. The greatest amount of CGRP release occurred in dental pulp from women in the luteal phase of the menstrual cycle. These results indicate that 5HT enhances capsaicin-evoked CGRP release from human trigeminal nociceptors in a sexually dimorphic manner providing a mechanistic basis for prevalence of trigeminal pain disorders in women.

Published by Elsevier B.V. on behalf of International Association for the Study of Pain.

### 1. Introduction

As approximately 1 in 4 persons experiences persistent craniofacial pain mediated by the trigeminal system, such as toothache and temporomandibular joint disorder pain [30,31] and approximately 10 million Americans experience migraine [40]. Craniofacial pain represents a prevalent and problematic burden, especially to the higher proportion of patients with improperly controlled pain [6,19]. Interestingly, trigeminal pain is more prevalent in women compared with men. For instance, migraine is twice as prevalent in women [29,30,42] and varies across the menstrual cycle [41,46]. However, it is unclear whether this sex difference in prevalence is due to biological differences, psychosocial differences, environmental differences, or to multiple factors.

The monoamine neurotransmitter serotonin (5HT) has been implicated in trigeminal pain, including migraine and masseter muscle pain [13,14,20,23], and 5HT is known to evoke hyperalgesia when injected into human tissues [4,13,14]. Therapeutics targeting the 5HT system have proved successful in treating some forms of trigeminal pain [2,10,16,38]. However, it remains unclear how 5HT modulates human trigeminal nociceptors. Preclinical animal models have provided insight into the mechanisms by which the trigeminal system mediates pain, and indicate that 5HT may regulate a subpopulation of trigeminal sensory neurons that express the transient receptor potential (TRPV1) ion channel, which is critical in transducing many noxious stimuli. TRPV1 is gated by thermal stimuli [7,8,12,45], oxidized linoleic acid metabolites [34,35,37], and inflammatory mediators [9,36] to induce calcium influx in nociceptors, resulting in release of inflammatory peptides such as calcitonin gene-related peptide (CGRP). There is evidence in dorsal root ganglia sensory neurons that 5HT potentiates TRPV1 functions [24,33,39] and prolongs nociceptor excitation during inflammation [1,22]. We recently reported that 5HT evokes calcium influx in

\* Corresponding author. Address: Pain Management Research Area, US Army Institute of Surgical Research, 3698 Chambers Pass, Fort Sam Houston, TX 78234, USA. Tel.: +1 (210) 539 9338; fax: +1 (210) 539 1460.

E-mail address: dayna.laveritt@us.army.mil (D.R. Loyd).

**Table 1**  
Demographic data of dental pulp donors.

Sex	Donors (n)	Teeth (n)	Age, y Mean (range)	Ethnicity
Male	22	45	22 (15–31)	23% Hispanic (n = 5) 55% White non-Hispanic (n = 12) 14% African American (n = 3) 4% Japanese/Asian (n = 1) 4% Persian/Indian (n = 1)
Female	33	95	22 (16–31)	39% Hispanic (n = 13) 52% White non-Hispanic (n = 17) 9% Japanese/Asian (n = 3)

capsaicin-sensitive rat trigeminal sensory neurons with a corresponding enhancement of CGRP release [26], and that 5HT<sub>2A</sub> and 5HT<sub>3A</sub> receptor antagonists and the anti-migraine drug sumatriptan (5HT<sub>1B/1D</sub> agonist), attenuate 5HT enhancement of capsaicin-evoked CGRP release *in vitro* and thermal hyperalgesia *in vivo* [25]. However, due to limitations of readily available human tissues, translation of these mechanistic findings into clinical relevance has been challenging.

Human dental pulp is composed of many of the same cells, fibers, and nociceptors as other human tissues, thus offering a readily available model for the study of human nociceptors [21]. Nociceptors innervating human dental pulp express TRPV1 and contain the nociceptive neuropeptide CGRP [15]. Using a recently developed *in vitro* superfusion method to measure CGRP release from human dental pulp [15], we tested our hypothesis that 5HT enhances capsaicin-evoked CGRP release from human nociceptors. We also performed Western blotting in human dental pulp to analyze the expression of 5HT receptors known to be involved in peripheral pain processing (5HT<sub>1B</sub>, 5HT<sub>1D</sub>, 5HT<sub>2A</sub>, and 5HT<sub>3A</sub>) with the aim of obtaining insight into the ability of 5HT to modulate human nociceptors.


## 2. Methods and materials

### 2.1. Study subjects

A total of 55 patients (22 male and 33 non-pregnant, post-pubertal, pre-menopausal female individuals) between the ages of 14 and 40 years who presented to the clinics of the University of Texas Health Science Center at San Antonio School of Dentistry consented to participate in this study. A total of 140 teeth were collected from enrolled patients who had already elected to have 1 to 4 third molars (“wisdom teeth”) extracted. Selected teeth were limited to a clinical diagnosis of a normal pulp lacking caries or restorations with no periapical radiolucencies upon radiographic evaluation and normal responsiveness to sensory testing. All teeth used in this study were required to show fully developed roots to ensure that the innervation of the pulp was complete. All patients were anesthetized using nerve block injection consisting of 2% lidocaine with epinephrine, with 78% of patients also receiving intravenous fentanyl and midazolam (50–100 µg and 5 mg, respectively) during extractions. Patient data collected for analysis included gender, age, ethnicity (Table 1), current medications, hormone-releasing birth control use, and days since last menses. All studies were approved by the University of Texas Health Science Center at San Antonio Institutional Review Board, and patients provided written informed consent to participate.

### 2.2. Materials

Capsaicin (TRPV1 agonist; Sigma, St. Louis, MO) was dissolved in ethanol stock and stored in aliquots at –20°C. A fresh aliquot was diluted for each experiment, with final dilutions containing




**Fig. 1.** Capsaicin evokes significant calcitonin gene-related peptide (CGRP) release from human dental pulp. Capsaicin (1–30 µmol/L; filled bars) induces CGRP release in a concentration-dependent manner from human dental pulp compared with vehicle (open bars), as expressed by amount (fmol) released per pulp (A). The same data are expressed as percentage of basal release (B). \*Significant at  $P < .05$  compared with vehicle;  $n = 15$ –18 teeth per group.

<0.5% ethanol. Serotonin hydrochloride (5HT; Sigma, St. Louis, MO) was dissolved in double-distilled water and diluted immediately before each use. All experimental vehicles were made at the same time and of the same constitution as the experimental drug.

### 2.3. *In vitro* superfusion of human dental pulp

Within 30 minutes of extraction, teeth were bisected and coronal pulp tissue was removed and placed into wells containing Hanks buffered saline solution (HBSS), pH 7.4 (Invitrogen, San Diego, CA). After an *in vitro* superfusion technique previously described [15], capsaicin responsiveness was determined by incubating the pulpal tissue (1 sample per well) in HBSS buffer alone for 20 minutes (basal sample) followed by a 20-minute exposure to vehicle or capsaicin (1 µmol/L, 10 µmol/L, or 30 µmol/L;  $n = 15$ –18 per group). To determine the effect of pretreatment with 5HT, pulpal tissue was incubated for 20 minutes in vehicle or 5HT



**Fig. 2.** Serotonin (5HT) significantly enhances calcitonin gene-related peptide (CGRP) release in female, but not male, human dental pulp. Serotonin (5HT) (100  $\mu\text{mol/L}$ ; filled bars;  $n = 35$ ) compared with vehicle (open bars;  $n = 32$ ) does not have an effect on CGRP release alone, but does significantly enhance capsaicin 1- $\mu\text{mol/L}$ -evoked CGRP release from tooth pulp (males and females combined) (A). When stratified by sex, there is no effect of 5HT on capsaicin-evoked CGRP release from male dental pulp ( $n = 17$ ) compared with vehicle ( $n = 12$ ; B). 5HT ( $n = 18$ ) significantly enhanced capsaicin-evoked CGRP release from dental pulp of females compared with vehicle ( $n = 20$ ) (C). Capsaicin or vehicle alone induced a comparable release of CGRP from the dental pulp of males (open bars,  $n = 12$ ) and females (filled bars;  $n = 20$ ) (D). \*Significant at  $P < .05$  compared with vehicle.

(100  $\mu\text{mol/L}$ ), a concentration shown to significantly enhance capsaicin-stimulated CGRP release in sensory neuron cultures [26], before stimulation with capsaicin (1  $\mu\text{mol/L}$ ;  $n = 32$ –35 per group). An additional group of 48 teeth from female patients was added to this group to conduct secondary analysis of the effects of hormone-releasing birth control use or menstrual cycle status on CGRP release. Upon completion of the superfusion experiments, tissue samples were lysed to release total cellular stores of CGRP by freezing/thawing the samples. Each superfusate sample (baseline, pretreatment, and capsaicin stimulated) was analyzed for CGRP levels by human-specific CGRP enzyme-linked immunoassay (SPI-BIO, Montigny le Bretonneux, France). Data are expressed as percentage of basal release to account for individual pulp variability and to avoid artificial inflation.

#### 2.4. Western blot

Protein was extracted from the dental pulp of 7 male and 6 female human third molar teeth. Teeth were cut on a sagittal plane and the pulp was extracted. The pulp was quickly placed in ice-cold phosphate-buffered saline solution (PBS) and then transferred into ice-cold homogenization buffer. Tissue homogenization was performed at 50 oscillations per second using a Tissue Lyzer LT (Qiagen, Valencia, CA) and a 5-mmol/L stainless steel bead. The pulp/homogenization buffer solution was lysed in 2-minute intervals, for a total of 4 minutes, to maintain the cool temperature of solution. Protein was then quantified using

a bicinchoninic acid assay. Equal amounts of protein were loaded and separated by polyacrylamide gel electrophoresis on a 12% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) gel and transferred to a polyvinylidene difluoride (PVDF) membrane (Bio-Rad, Hercules, CA). The membrane was then blocked with 5% milk in TBS-Tween 20, and 5HT receptors were probed with selective antibodies for either goat anti-5HT<sub>1B</sub> receptor (1:500; 42 kDa; Santa Cruz Biotechnology, Santa Cruz, CA), goat anti-5HT<sub>1D</sub> receptor (1:500; 43 kDa; Santa Cruz Biotechnology), rabbit anti-5HT<sub>2A</sub> receptor (1:500; 53 kDa; Immunostar, Hudson, WI) or rabbit anti-5HT<sub>3A</sub> receptor (1:1000; 53 kDa; Proteintech, Chicago, IL) and monoclonal mouse anti- $\beta$ -actin (1:1000; Santa Cruz Biotechnology) as control. Membranes were then treated with either donkey anti-goat (1:2500; Santa Cruz Biotechnology), goat anti-rabbit (1:2500 or 1:5000) or goat anti-mouse (1:5000; Santa Cruz Biotechnology) secondary antisera linked to horseradish peroxidase for signal amplification and detected with an enhanced chemiluminescence reagent kit (Amersham, Piscataway, NJ). To confirm antibody selectivity, blots were pre-incubated in blocking peptides for the 5HT<sub>1B</sub> (1:500; Santa Cruz Biotechnology), 5HT<sub>1D</sub> (1:500; Santa Cruz Biotechnology) and 5HT<sub>2A</sub> (1:500; Immunostar). A blocking peptide was unavailable for the 5HT<sub>3A</sub> antibody; a lack of staining was confirmed by omitting the primary antibody. Images were captured with an AlphaImager Gel Documentation system (Cell Biosciences, Santa Clara, CA), and brightness/contrast values were maximized for publication with Adobe Photoshop CS4.

## 2.5. Data analysis

Superfusion data are presented as mean  $\pm$  SEM percentage of basal levels. Western blots were analyzed by densitometry using ImageJ-64 (<http://www.rsb.info.nih.gov/ij/>). Each blot was selected as a region of interest, sampled 3 times, and analyzed for average gray scale pixel value (sum gray values/number of pixels; 16-bit) following background correction (set at  $\sim$ 150 pixels) [17]. Data are reported as percentage of  $\beta$ -actin controls. Statistical analyses using GraphPad Prism software version 5 (GraphPad, San Diego, CA) were conducted by unpaired t-test, or one-way or 2-way analysis of variance and *post hoc* individual group analyses were compared using the Bonferroni *post hoc* test. Statistical significance was tested at  $P < .05$ . Error bars represent the standard error of the mean.

## 3. Results

### 3.1. Capsaicin evokes significant, concentration-dependent CGRP release from human dental pulp


Capsaicin alone evoked a concentration-dependent increase in CGRP release ranging from  $28 \pm 3$  to  $119 \pm 12$  fmol/pulp at 1 to 30  $\mu$ mol/L compared with vehicle controls (Fig. 1A). When analyzed as percentage of basal CGRP release per pulp, this increase in CGRP release was significant across all concentrations compared with vehicle ( $F_{2,92} = 4.04$ ;  $P < .05$ ), with the 10- and 30- $\mu$ mol/L concentration more than doubling CGRP release (Fig. 1B). The 10- and 30- $\mu$ mol/L concentrations produced significantly greater degree of CGRP release compared with the 1- $\mu$ mol/L concentration. The 1- $\mu$ mol/L concentration was chosen for the next set of experiments to permit detection of a potential enhancement of capsaicin-evoked CGRP release.

### 3.2. Serotonin enhances capsaicin-stimulated CGRP release from female but not male human dental pulp

Pretreatment with 5HT 100  $\mu$ mol/L significantly increased capsaicin-stimulated CGRP release by  $\sim$ 50% ( $F_{1,134} = 14.88$ ;  $P < .05$ ), whereas 5HT alone did not evoke CGRP release ( $P > .05$ ) (Fig. 2A). When these data were stratified by sex, there was no significant effect of 5HT on capsaicin-stimulated CGRP release from male dental pulp ( $F_{1,54} = 1.18$ ; NS) (Fig. 2B). However, 5HT pretreatment evoked a significant, 2-fold enhancement of CGRP release in female dental pulp ( $F_{1,72} = 20.15$ ;  $P < .05$ ) (Fig. 2C). This effect was independent of whether the tooth was erupted or non-erupted before extraction [ $t(16) = 1.53$ ; NS]. The observed sexually dimorphic effect of 5HT-enhancement of capsaicin-evoked CGRP release in human dental pulp may be due to differences in CGRP concentrations or TRPV1 levels. To evaluate these possibilities, we administered capsaicin alone to male and female dental pulp and measured evoked CGRP release. Capsaicin alone did not evoke a significantly different level of CGRP release from male and female dental pulp ( $F_{1,60} = 0.05$ ; NS) (Fig. 2D).

### 3.3. Serotonin receptors are expressed at similar levels in male and female human dental pulp

The expression of 5HT receptors in human dental pulp was examined by Western blot to confirm the availability of anatomical targets for 5HT modulation. The 5HT<sub>1B</sub>, 5HT<sub>1D</sub>, 5HT<sub>2A</sub>, and 5HT<sub>3A</sub> receptor protein was expressed in both male and female human dental pulp (Fig. 3A). Densitometry on the western blots revealed no significant differences in the expression of these receptor



**Fig. 3.** Serotonin (5HT) receptor protein is expressed in human dental pulp and is comparable between the sexes. Western blot bands illustrating 5HT<sub>1B</sub>, 5HT<sub>1D</sub>, 5HT<sub>2A</sub>, and 5HT<sub>3A</sub> receptor expression in male versus female human dental pulp (A). Quantification of Western blots indicates no sex differences in expression; male,  $n = 7$ ; female,  $n = 6$  (B).


subtypes between male and female dental pulp [ $t(11) = 0.69, 0.86, 1.12, \text{ and } 0.18$ , respectively; NS] (Fig. 3B).

### 3.4. Effect of age and ethnicity on 5HT enhancement of capsaicin-stimulated CGRP release from male and female human dental pulp

5HT produced consistently higher capsaicin-evoked CGRP release from dental pulp of female individuals who were more than 24 years of age compared with those 15 to 20 years of age ( $F_{3,17} = 5.76$ ;  $P < .05$ ) (Fig. 4A). There was no effect of age on capsaicin-stimulated CGRP release after vehicle pretreatment ( $F_{5,19} = 2.886$ ; NS), and there was no significant effect of age on dental pulp from males ( $F_{5,16} = 0.58$ ; NS) (Fig. 4B). The vast majority of dental pulp was from non-Hispanic white and Hispanic patients. When stratified by these 2 groups, there was no significant effect of ethnicity on 5HT enhancement of CGRP release from female ( $F_{1,34} = 3.52$ ; NS) (Fig. 4C) or male ( $F_{1,16} = 0.27$ ; NS) dental pulp (Fig. 4D).

### 3.5. Female patients who were amenstrual because of hormonal IUD or were in the week before menses presented with the greatest levels of 5HT-enhanced CGRP release

The CGRP release data from the dental pulp of female patients were divided into 3 groups based on these individuals' current use



**Fig. 4.** There is a significant effect of age, but not ethnicity, on serotonin (5HT) enhancement of calcitonin gene-related peptide (CGRP) release from female dental pulp. Effect of 5HT on capsaicin-evoked CGRP release stratified by age in females (A;  $n = 18$ ) versus males (B;  $n = 17$ ) and by ethnicity in females (C;  $n = 8$  white non-Hispanic and 10 Hispanic) versus males (D;  $n = 10$  white non-Hispanic and 7 Hispanic). \*Significant at  $P < .05$  compared with vehicle.

of hormonal manipulations to prevent pregnancy: use of oral birth control (OBC), no use of oral birth control (No OBC), or amenstrual because of the use of a synthetic progestogen-releasing intrauterine device (Amenstrual/IUD). 5HT-enhanced CGRP release was significantly higher in dental pulp from female subjects who were amenstrual because of an IUD ( $F_{2,52} = 14.92$ ;  $P < .05$ ) (Fig. 5A). 5HT-enhanced CGRP release was similar in the dental pulp of females regardless of the use of oral birth control. The CGRP release data from the dental pulp of female patients not using hormonal manipulations (No OBC) was then further divided into 4 groups based on the first day of the last menses: days 1 to 7, 8 to 14, 15 to 21, and 22 to 28 days. There was a significant effect of the status of menstrual cycle on 5HT enhancement of capsaicin-stimulated CGRP release ( $F_{3,41} = 3.06$ ;  $P < .05$ ) (Fig. 5B). 5HT-enhanced CGRP release was lowest in dental pulp from females in the week during menses (days 1–7), whereas 5HT-enhanced CGRP release was highest in dental pulp from females in the week before menses (days 22–28). In contrast, there was no significant effect of day since last menses on CGRP release evoked by capsaicin alone ( $F_{3,18} = 1.714$ ; NS).

#### 4. Discussion


5HT is a pronociceptive mediator in the periphery that has been reported to enhance TRPV1-evoked CGRP release from rat trigeminal sensory neurons [26]; however, whether this occurs in human nociceptors is unknown. Using an *in vitro* superfusion assay on human dental pulp, here we report that (1) 5HT enhances capsaicin-evoked CGRP release from female dental pulp, but not male dental pulp; and (2) that 5HT-enhanced CGRP release varies across age and the menstrual cycle. Furthermore, we report that there are no significant sex differences in 5HT<sub>1B</sub>, 5HT<sub>1D</sub>, 5HT<sub>2A</sub>, or 5HT<sub>3A</sub>

receptor protein expression in human dental pulp, and that capsaicin evokes similar levels of CGRP release from both male and female peptidergic terminals.

Capsaicin steadily evoked a concentration-dependent increase in CGRP release consistent with that previously reported [15]. CGRP release to vehicle treatment was consistently lower than basal levels, likely because of further stabilization of the extracted dental pulp. Capsaicin produces maximal CGRP release at 60  $\mu\text{mol/L}$  without inducing detectable desensitization, which is observed at 100 and 300  $\mu\text{mol/L}$  in dental pulp [15]. In this study, capsaicin 1 to 30  $\mu\text{mol/L}$  was examined to allow for potential enhancement of capsaicin-evoked CGRP release without inducing desensitization. Similar to previous studies [15], there was no detectable effect of local nerve block in the absence or presence of intravenous fentanyl/midazolam on CGRP release from human dental pulp. As capsaicin 1  $\mu\text{mol/L}$  was the lowest concentration to evoke a significant increase in CGRP release when analyzed as percentage of basal release, this concentration was chosen to stimulate CGRP release after pretreatment with either 5HT or vehicle. When dental pulp was pretreated with 5HT before capsaicin, capsaicin-evoked CGRP release was significantly enhanced. Importantly, 5HT alone did not evoke a significant change in CGRP release. Together, these data indicate that 5HT modulates TRPV1-evoked CGRP release in human dental pulp. This modulation may be occurring via direct or indirect actions on TRPV1-expressing nociceptors in human dental pulp. 5HT receptors may be located on non-neuronal cells; however, it is unlikely that 5HT modulates CGRP release through non-TRPV1-expressing cells, as 5HT given alone did not alter CGRP release.

As trigeminal pain is more prevalent in women, the present data were then stratified by sex of the dental pulp donor to determine potential sex differences in 5HT-evoked CGRP release.





**Fig. 5.** Serotonin (5HT) enhanced calcitonin gene-related peptide (CGRP) release to a greater degree in dental pulp from female patients during the week before menses. Effect of 5HT on capsaicin-evoked CGRP release grouped by teeth from females not on oral birth control (No OBC), on oral birth control (OBC), or amenstrual because of hormone-releasing IUD implantation (Amenstrual/IUD; A). Data from patients not on birth control are also expressed as days since first day of last menstrual cycle (B). \*Significant at  $P < .05$  compared with vehicle.

Indeed, 5HT pretreatment enhanced capsaicin-stimulated CGRP release from the dental pulp of females only. There was no significant effect of 5HT pretreatment on the dental pulp from males. It is important to note that capsaicin evoked comparable CGRP release from male and female dental pulp, indicating that the observed effect is due to differences in serotonergic sensitization of TRPV1 neurons. As there were no sex differences in capsaicin-evoked or 5HT-evoked CGRP release, and as there was no difference in the 5HT receptor proteins analyzed, a remaining possibility is that 5HT receptor coexpression on TRPV1 fibers is sexually dimorphic. Because there are current limitations on available specific 5HT receptor antibodies, this possibility remains untested. There was also no effect of whether the donors were of non-Hispanic white or Hispanic ethnicity on CGRP release. Together, these data provide evidence of a sexually dimorphic biological mechanism that may contribute to the well-recognized sex differences in some forms of trigeminal pain. Although it has been reported that 5HT-evoked rat trigeminal afferent discharge is not sexually dimorphic [44], our data indicate that 5HT may instead alter pain transmission indirectly via enhancing TRPV1-evoked CGRP release.

It is possible that the observed sexually dimorphic serotonin effect is mediated by the actions of sex steroids. Estradiol potentiates capsaicin-mediated currents in rat dorsal root ganglia sensory neurons [11], and female rodents exhibit greater capsaicin-evoked nocifensive responses during proestrous [27], when estrogen levels peak. To examine whether 5HT-enhanced CGRP release is altered across the menstrual cycle, we stratified our data first by whether

the donor was on oral or IUD-mediated birth control. Interestingly, dental pulp from patients who were amenstrual because of a hormone-releasing IUD [28] represented the highest levels of 5HT-enhanced CGRP release. The data from female individuals on oral birth control, and approximately half of those not on oral birth control, were similar to CGRP release observed in male pulp (~170%), whereas the data from amenstrual females was ~300%; thus, when the data are combined, the amount of CGRP release is ~230% (Fig. 2). When the data from normally cycling female subjects (No OBC) are stratified by days since last menses, dental pulp from those in the week before menses evoked the highest amount of 5HT-enhanced CGRP release. These data are interesting, as both estrogen and, particularly, progesterone significantly increase during the luteal phase (days 16–28) of the menstrual cycle during the week before menses [43].

Importantly, there was no significant effect of day since last menses on CGRP release evoked by capsaicin alone, providing further support for a hypothesis that sex steroids specifically affect 5HT modulation of TRPV1 nociceptors. There was an effect of age on 5HT-enhanced CGRP release in female dental pulp, whereas there was no effect of age on capsaicin-evoked CGRP release or CGRP release in male dental pulp, which could also indicate an effect of hormonal status in 5HT-enhancement of CGRP release. Our previous studies examining the enhancing effect of 5HT on TRPV1-evoked CGRP release and thermal hyperalgesia were limited to male rats, so whether this effect is observed in female rats is unknown. Preclinical studies examining the effects of 5HT and steroid hormones in female rats across the estrous cycle are warranted to address this possibility.

We also found that 4 different 5HT receptor subtypes known to be involved in 5HT-evoked pain processing [26,27], namely, 5HT<sub>1B</sub>, 5HT<sub>1D</sub>, 5HT<sub>2A</sub>, and 5HT<sub>3A</sub> receptors, were expressed in male and female human dental pulp. These data provide possible pharmacological targets by which the enhancing effects of 5HT on TRPV1-evoked CGRP release may be controlled. This is important, as 5HT receptor expression in the trigeminal system represents a critical target for reducing CGRP release [3,26], which is correlated with headache and migraine in humans. When quantified, the protein expression of these receptors was comparable between male and female dental pulp. Given our observed alterations of 5HT-enhanced CGRP release over the menstrual cycle, further studies are required to determine whether 5HT receptor expression is also altered across the menstrual cycle in human dental pulp. This possibility may be unlikely, given that 5HT<sub>1</sub> receptor mRNA levels in the mouse trigeminal ganglia do not fluctuate over the estrous cycle [5]; however, this may represent an effect that occurs in human but not rodent tissues and should be considered, or may not reflect changes in translational control.

Clinical evidence suggests that at least 1 form of trigeminal pain, headaches and migraine, fluctuates with menstrual cycle status. Headache and migraine typically occur in women around menses, and some women experience migraine only in association with menses [32]. Considering our data in this population, 5HT may be enhancing CGRP release from the TRPV1 population of trigeminal nociceptors at the onset of menses. Future studies examining whether this effect occurs via TRPV1 and/or via specific 5HT receptors would offer therapeutic insight and are warranted. Importantly, these data illustrate the necessity of examining both male and female subjects in studies of trigeminal pain [18]. Overall, these results indicate that 5HT enhances TRPV1-evoked CGRP release from female human dental pulp and provide evidence of a sexually dimorphic peripheral mechanism modulating trigeminal pain processing. Future studies examining serotonergics that block or attenuate 5HT enhancement of TRPV1-evoked CGRP release may prove therapeutic for elusive trigeminal pain disorders in women.

## Conflict of interest statement

The authors report no conflicts of interest.

## Acknowledgments

The authors would like to acknowledge the helpful comments of Michael A. Henry, DDS, PhD, and the technical assistance of Paul Chen. This work was supported by National Institutes of Health grants NCATS UL1TR000149 and R01 NS58655 (K.M.H.), T32 DE14318, F32 DE021309 (D.R.L.).

## References

- [1] Aley KO, Messing RO, Mochly-Rosen D, Levine JD. Chronic hypersensitivity for inflammatory nociceptor sensitization mediated by the epsilon isozyme of protein kinase C. *J Neurosci* 2000;20:4680–5.
- [2] Ali Z, Prabhakar H, Rath GP, Dash HH. Enoxaparin induced intracerebral haemorrhage after deep brain stimulation surgery. *Eur J Anaesthesiol* 2009;26:617–8.
- [3] Amrutkar DV, Ploug KB, Hay-Schmidt A, Porreca F, Olesen J, Jansen-Olesen I. MRNA expression of 5-hydroxytryptamine 1B, 1D, and 1F receptors and their role in controlling the release of calcitonin gene-related peptide in the rat trigeminovascular system. *PAIN<sup>®</sup>* 2012;153:830–8.
- [4] Babenko V, Svensson P, Graven-Nielsen T, Drewes AM, Jensen TS, Arendt-Nielsen L. Duration and distribution of experimental muscle hyperalgesia in humans following combined infusions of serotonin and bradykinin. *Brain Res* 2000;853:275–81.
- [5] Berman NE, Puri V, Chandrala S, Puri S, Macgregor R, Liverman CS, Klein RM. Serotonin in trigeminal ganglia of female rodents: relevance to menstrual migraine. *Headache* 2006;46:1230–45.
- [6] Bigal ME, Serrano D, Reed M, Lipton RB. Chronic migraine in the population: burden, diagnosis, and satisfaction with treatment. *Neurology* 2008;71:559–66.
- [7] Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, Koltzenburg M, Basbaum AI, Julius D. Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 2000;288:306–13.
- [8] Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, Julius D. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 1997;389:816–24.
- [9] Cesare P, Moriondo A, Vellani V, McNaughton PA. Ion channels gated by heat. *Proc Natl Acad Sci USA* 1999;96:7658–63.
- [10] Chen LC, Ashcroft DM. Meta-analysis of the efficacy and safety of zolmitriptan in the acute treatment of migraine. *Headache* 2008;48:236–47.
- [11] Chen SC, Chang TJ, Wu FS. Competitive inhibition of the capsaicin receptor-mediated current by dehydroepiandrosterone in rat dorsal root ganglion neurons. *J Pharmacol Exp Ther* 2004;311:529–36.
- [12] Davis JB, Gray J, Gunthorpe MJ, Hatcher JP, Davey PT, Overend P, Harries MH, Latcham J, Clapham C, Atkinson K, Hughes SA, Rance K, Grau E, Harper AJ, Pugh PL, Rogers DC, Bingham S, Randall A, Sheardown SA. Vanilloid receptor-1 is essential for inflammatory thermal hyperalgesia. *Nature* 2000;405:183–7.
- [13] Ernberg M, Hedenberg-Magnusson B, Kurita H, Kopp S. Effects of local serotonin administration on pain and microcirculation in the human masseter muscle. *J Orofac Pain* 2006;20:241–8.
- [14] Ernberg M, Lundeberg T, Kopp S. Pain and allodynia/hyperalgesia induced by intramuscular injection of serotonin in patients with fibromyalgia and healthy individuals. *PAIN<sup>®</sup>* 2000;85:31–9.
- [15] Fehrenbacher JC, Sun XX, Locke EE, Henry MA, Hargreaves KM. Capsaicin-evoked iCGRP release from human dental pulp: a model system for the study of peripheral neuropeptide secretion in normal healthy tissue. *PAIN<sup>®</sup>* 2009;144:253–61.
- [16] Ferrari MD, Roon KI, Lipton RB, Goadsby PJ. Oral triptans (serotonin 5-HT<sub>1B/1D</sub>) agonists) in acute migraine treatment: a meta-analysis of 53 trials. *Lancet* 2001;358:1668–75.
- [17] Gassmann M, Grenacher B, Rohde B, Vogel J. Quantifying Western blots: pitfalls of densitometry. *Electrophoresis* 2009;30:1845–55.
- [18] Greenspan JD, Craft RM, LeResche L, Arendt-Nielsen L, Berkley KJ, Fillingim RB, Gold MS, Holdcroft A, Lautenbacher S, Mayer EA, Mogil JS, Murphy AZ, Traub RJ. Studying sex and gender differences in pain and analgesia: a consensus report. *PAIN<sup>®</sup>* 2007;132:S26–45.
- [19] Hargreaves KM. Orofacial pain. *PAIN<sup>®</sup>* 2011;152:S25–32.
- [20] Hargreaves RJ, Shephard SL. Pathophysiology of migraine—new insights. *Can J Neurol Sci* 1999;26:S12–9.
- [21] Henry MA, Hargreaves KM. Peripheral mechanisms of odontogenic pain. *Dent Clin North Am* 2007;51:19–44.
- [22] Herbert MK, Schmidt RF. Activation of normal and inflamed fine articular afferent units by serotonin. *PAIN<sup>®</sup>* 1992;50:79–88.
- [23] Herken H, Erdal E, Mutlu N, Barlas O, Cataloluk O, Oz F, Guray E. Possible association of temporomandibular joint pain and dysfunction with a polymorphism in the serotonin transporter gene. *Am J Orthod Dentofacial Orthop* 2001;120:308–13.
- [24] Hisaoka T, Morikawa Y, Kitamura T, Senba E. Expression of a member of tumor necrosis factor receptor superfamily, TROY, in the developing olfactory system. *Glia* 2004;45:313–24.
- [25] Loyd DR, Chen PB, Hargreaves KM. Anti-hyperalgesic effects of anti-serotonergic compounds on serotonin- and capsaicin-evoked thermal hyperalgesia in the rat. *Neuroscience* 2012;203:207–15.
- [26] Loyd DR, Weiss G, Henry MA, Hargreaves KM. Serotonin increases the functional activity of capsaicin-sensitive rat trigeminal nociceptors via peripheral serotonin receptors. *PAIN<sup>®</sup>* 2011;152:2267–76.
- [27] Lu YC, Chen CW, Wang SY, Wu FS. 17Beta-estradiol mediates the sex difference in capsaicin-induced nociception in rats. *J Pharmacol Exp Ther* 2009;331:1104–10.
- [28] Luukkainen T, Lahteenmaki P, Toivonen J. Levonorgestrel-releasing intrauterine device. *Ann Med* 1990;22:85–90.
- [29] Macfarlane TV, Blinkhorn AS, Davies RM, Kincey J, Worthington HV. Oro-facial pain in the community: prevalence and associated impact. *Commun Dent Oral Epidemiol* 2002;30:52–60.
- [30] Macfarlane TV, Blinkhorn AS, Davies RM, Kincey J, Worthington HV. Predictors of outcome for orofacial pain in the general population: a four-year follow-up study. *J Dent Res* 2004;83:712–7.
- [31] Macfarlane TV, Blinkhorn AS, Davies RM, Ryan P, Worthington HV, Macfarlane GJ. Orofacial pain: just another chronic pain? Results from a population-based survey. *PAIN<sup>®</sup>* 2002;99:453–8.
- [32] Marcus DA. Interrelationships of neurochemicals, estrogen, and recurring headache. *PAIN<sup>®</sup>* 1995;62:129–39.
- [33] Ohta T, Ikemi Y, Murakami M, Imagawa T, Otsuguro K, Ito S. Potentiation of transient receptor potential V1 functions by the activation of metabotropic 5-HT receptors in rat primary sensory neurons. *J Physiol* 2006;576:809–22.
- [34] Patwardhan AM, Akopian AN, Ruparel NB, Diogenes A, Weintraub ST, Uhlson C, Murphy RC, Hargreaves KM. Heat generates oxidized linoleic acid metabolites that activate TRPV1 and produce pain in rodents. *J Clin Invest* 2010;120:1617–26.
- [35] Patwardhan AM, Scotland PE, Akopian AN, Hargreaves KM. Activation of TRPV1 in the spinal cord by oxidized linoleic acid metabolites contributes to inflammatory hyperalgesia. *Proc Natl Acad Sci USA* 2009;106:18820–4.
- [36] Pingle SC, Matta JA, Ahern GP. Capsaicin receptor: TRPV1 a promiscuous TRP channel. *Handb Exp Pharmacol* 2007:155–71.
- [37] Ruparel S, Henry MH, Akopian A, Patil NB, Zeldin D, Roman L, Hargreaves KM. Plasticity of cytochrome P450 isozyme expression in rat trigeminal neurons during inflammation. *PAIN<sup>®</sup>* 2012;153:2031–9.
- [38] Schabitz WR, Schade H, Heiland S, Kollmar R, Bardutzky J, Henninger N, Muller H, Carl U, Toyokuni S, Sommer C, Schwab S. Neuroprotection by hyperbaric oxygenation after experimental focal cerebral ischemia monitored by MRI. *Stroke* 2004;35:1175–9.
- [39] Simonetti M, Fabbro A, D'Arco M, Zweyer M, Nistri A, Giniatullin R, Fabbretti E. Comparison of P2X and TRPV1 receptors in ganglia or primary culture of trigeminal neurons and their modulation by NGF or serotonin. *Mol Pain* 2006;2:11.
- [40] Stewart WF, Lipton RB, Celentano DD, Reed ML. Prevalence of migraine headache in the United States. Relation to age, income, race, and other sociodemographic factors. *JAMA* 1992;267:64–9.
- [41] Stewart WF, Lipton RB, Chee E, Sawyer J, Silberstein SD. Menstrual cycle and headache in a population sample of migraineurs. *Neurology* 2000;55:1517–23.
- [42] Stewart WF, Shechter A, Rasmussen BK. Migraine prevalence. A review of population-based studies. *Neurology* 1994;44:S17–23.
- [43] Stricker R, Eberhart R, Chevailler MC, Quinn FA, Bischof P. Establishment of detailed reference values for luteinizing hormone, follicle stimulating hormone, estradiol, and progesterone during different phases of the menstrual cycle on the Abbott ARCHITECT analyzer. *Clin Chem Lab Med* 2006;44:883–7.
- [44] Sung D, Dong X, Ernberg M, Kumar U, Cairns BE. Serotonin (5-HT) excites rat masticatory muscle afferent fibers through activation of peripheral 5-HT<sub>3</sub> receptors. *PAIN<sup>®</sup>* 2008;134:41–50.
- [45] Voets T, Droogmans G, Wissenbach U, Janssens A, Flockerzi V, Nilius B. The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nature* 2004;430:748–54.
- [46] Welch KM. Migraine and ovarian steroid hormones. *Cephalalgia* 1997;17:12–6.